

Anomalous Spin Dynamics of Hubbard Model on Honeycomb Lattices

Gao-Yong Sun¹ and Su-Peng Kou^{1,*}

¹*Department of Physics, Beijing Normal University, Beijing 100875, China*

In this paper, the honeycomb Hubbard model in optical lattices is investigated using O(3) non-linear σ model. A possible quantum non-magnetic insulator in a narrow parameter region is found near the metal-insulator transition. We study the corresponding dynamics of magnetic properties, and find that the narrow region could be widened by hole doping.

PACS numbers: 71.10.Fd, 71.10.Hf, 75.10.-b, 71.30.+h

I. INTRODUCTION

Recently, using ultracold atoms to form Bose-Einstein Condensates (BEC) or Fermi degenerate gases for precise measurements and simulations of quantum many-body systems, is quite impressive and has become a rapidly-developing field^{1,2}. Experimental realizations of quantum many-body systems in optical lattices have made a chance to simulate strongly correlated systems¹. People have successfully observed the Mott insulator–superfluid transition both in bosonic³ and fermionic atoms (e.g. ⁶Li or ⁴⁰K, etc.)^{4,5}. In particular, the two-dimensional (fermionic) Hubbard model is one of the most interesting issues depicting the nature of high-temperature superconductivity. It is still a big challenge to clearly understand the physics of repulsive Hubbard model on two dimensional (2D) lattices. Thus people try to simulate the Fermi-Hubbard model using a two-component mixture of repulsively interacting fermions^{6,7,8,9}.

In this paper, we focus on the two dimensional Hubbard model in honeycomb lattices (2D honeycomb Hubbard model). In Refs.^{10,11,12}, it is proposed that the 2D honeycomb optical lattice can be realized in the cold atoms with three detuned standing-wave lasers, of which the optical potential is given by

$$V(x, y) = \sum_{j=1,2,3} V \sin^2[k_L(x \cos \theta_j + y \sin \theta_j) + \pi/2] \quad (1)$$

where $\theta_1 = \pi/3$, $\theta_2 = 2\pi/3$, $\theta_3 = 0$, and k_L is the optical wave vector. When two-component fermions with repulsive interaction are put into the 2D honeycomb optical lattice, one can get an effective honeycomb Hubbard model. It is easy to change the potential barrier V by varying the laser intensities to tune the Hamiltonian parameters including the hopping strength t (t -term) and the particle interaction U (U -term). This lays the foundation for our later discussions and calculations.

The simplest approach to study the honeycomb Hubbard model is the Hartree-Fock (HF) mean field method, from which people obtain a semi-metal-insulator (MI) transition at a critical value $(U/t)_c$ between a semi-metal (SM) and an antiferromagnetic (AF) insulator. In the weak interaction region $U/t < (U/t)_c$, the ground state is a semi-metal (SM) with nodal fermi-points. In the strong interaction region $U/t > (U/t)_c$, the ground state becomes an insulator with massive fermionic excitations. However, the HF theory does not keep the spin rotation




FIG. 1: Illustration of a honeycomb lattice with A and B sublattices. a is the length of the hexagon side and chosen to be unit.

symmetry by fixing the spins along $\hat{z}$ -axis. So the results are not reliable. Though the semi-metal-insulator transition of the honeycomb Hubbard model has been studied by different approaches,^{13,14,15,16} the results are not consistent with each other.

In this paper, we will investigate the two dimensional honeycomb Hubbard model by an approach proposed in Refs.^{17,18,19,20,21,22} that keeps spin rotation symmetry. By it, we find anomalous spin dynamics not far from the critical point of MI transition : there may exist a narrow non-magnetic insulator. The narrow non-magnetic insulator will be in favor of hole-doping.

The paper is organized as follows. In Sec. II, the semi-metal-insulator transition is studied by HF mean field approach. In Sec. III, an effective O(3) non-linear σ model (NL σ M) is obtained to investigate properties of the honeycomb-Hubbard model. In Sec. IV, a global phase diagram is given and magnetic properties of the insulator state is studied based on the NL σ M. In Sec. V, we discuss how to observe the anomalous spin dynamics in optical lattice of cold atoms. In Sec. VI, we show the doping effect on the magnetic properties of the ground state. Finally, the conclusions are given in Sec. VII.

II. METAL-INSULATOR TRANSITION

As a starting point, the Hamiltonian of the Hubbard model on honeycomb lattice is given by

$$\mathcal{H} = -t \sum_{\langle i,j \rangle} (\hat{c}_i^\dagger \hat{c}_j + h.c.) + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} - \mu \sum_i \hat{c}_i^\dagger \hat{c}_i. \quad (2)$$

Here $\hat{c}_i = (\hat{c}_{i\uparrow}, \hat{c}_{i\downarrow})^T$ and $\hat{c}_{i,\sigma}^\dagger, \hat{c}_{i,\sigma}$ are electronic creation and annihilation operators. t is the hopping integral. U is the on-site Coulomb repulsion. σ are the spin-indices representing spin-up ($\sigma = \uparrow$) and spin-down ($\sigma = \downarrow$) for electrons. μ is the chemical potential which is $\frac{U}{2}$ in half-filling. $\langle i, j \rangle$ denotes two sites on a nearest-neighbor link. $\hat{n}_{i\uparrow}$ and $\hat{n}_{i\downarrow}$ are the number operators of electrons with up-spin and down spin respectively.

Because the honeycomb lattice is a bipartite lattice (See Fig.1), we divide the system into two sublattices, A and B. Using the Fourier transformations, the electronic annihilation operators on two sublattices are written into

$$\hat{c}_{i \in A, \sigma} = \frac{1}{\sqrt{N_s}} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{R}_i} \hat{a}_{\mathbf{k}\sigma}, \quad (3)$$

$$\hat{c}_{i \in B, \sigma} = \frac{1}{\sqrt{N_s}} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{R}_i} \hat{b}_{\mathbf{k}\sigma}, \quad (4)$$

where N_s denoting the number of unit cells. For free fermions, the Hamiltonian could be transformed in the momentum space as

$$\mathcal{H} = \sum_{\mathbf{k}, \sigma} \begin{pmatrix} \hat{a}_{\mathbf{k}\sigma}^\dagger & \hat{b}_{\mathbf{k}\sigma}^\dagger \end{pmatrix} \begin{pmatrix} 0 & \xi_{\mathbf{k}} \\ \xi_{\mathbf{k}}^* & 0 \end{pmatrix} \begin{pmatrix} \hat{a}_{\mathbf{k}\sigma} \\ \hat{b}_{\mathbf{k}\sigma} \end{pmatrix} \quad (5)$$

where the energy of free fermions is

$$|\xi_{\mathbf{k}}| = \left| -t \sum_{\delta} e^{i\mathbf{k} \cdot \delta} \right| \quad (6)$$

$$= t \sqrt{3 + 2 \cos(\sqrt{3}k_y) + 4 \cos(3k_x/2) \cos(\sqrt{3}k_y/2)}.$$

Here the nearest neighbors of a electron in A lattice are defined as

$$\delta_1 = \frac{a}{2} (1, \sqrt{3}), \delta_2 = \frac{a}{2} (1, -\sqrt{3}), \delta_3 = (-a, 0) \quad (7)$$

where a is the length of the hexagon side and chosen to be unit. The spectrum for free electrons is then obtained as $E_{\mathbf{k}} = \pm |\xi_{\mathbf{k}}|$.

Next we use the path-integral formulation of electrons with spin rotation symmetry to study the on-site repulsive interaction in Hubbard model.^{17,18,19,20,21,22} The interaction term can be handled by using a SU(2) invariant Hubbard-Stratonovich (HS) decomposition in the arbitrary on-site unit vector $\mathbf{\Omega}_i$

$$\hat{n}_{i\uparrow} \hat{n}_{i\downarrow} = \frac{(\hat{c}_i^\dagger \hat{c}_i)^2}{4} - \frac{1}{4} [\mathbf{\Omega}_i \cdot \hat{c}_i^\dagger \sigma \hat{c}_i]^2 \quad (8)$$

where $\sigma = (\sigma_x, \sigma_y, \sigma_z)$ is the Pauli matrix. Then the HS transformation for the interaction term is

$$e^{U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow}} = \int \prod_i \frac{d\Delta_c d\Delta_i d^2 \mathbf{\Omega}_i}{4\pi^2 U} \exp \left(\sum_i \left[\frac{1}{U} (\Delta_c^2 + \Delta_i^2) + i\Delta_c \hat{c}_i^\dagger \hat{c}_i - \Delta_i \hat{c}_i^\dagger \mathbf{\Omega}_i \cdot \sigma \hat{c}_i \right] \right). \quad (9)$$

Here Δ_c and Δ_i are the auxiliary fields. By replacing electronic operators $\hat{c}_{i,\sigma}^\dagger$ and $\hat{c}_{i,\sigma}$ to Grassmann variables $c_{i,\sigma}^*$ and $c_{i,\sigma}$, the effective Lagrangian in terms of Grassmann variables $c_{i,\sigma}^*$ and $c_{i,\sigma}$ is then obtained as

$$\mathcal{L}_{\text{eff}} = \sum_{i,\sigma} c_{i,\sigma}^* \partial_\tau c_{i,\sigma} - \sum_{\langle ij \rangle} (t_{i,j} c_i^* c_j + h.c.) - \sum_i \Delta_i c_i^* \mathbf{\Omega}_i \cdot \sigma c_i + \sum_i \left[\frac{1}{U} (\Delta_c^2 + \Delta_i^2) + (i\Delta_c - \mu) c_i^* c_i \right]. \quad (10)$$

The ground state of the honeycomb Hubbard model is known to be long-range AF order in the large U limit. Such **an** order can be described by a simple saddle-point Lagrangian by fixing the direction vector field $\mathbf{\Omega}_i$ to $\hat{\mathbf{z}}$ -axis $\mathbf{\Omega}_i = (-1)^i \hat{\mathbf{z}}$ and choosing the amplitude as

$$i\Delta_c = \frac{U}{2} \langle c_i^\dagger c_i \rangle = \frac{Un}{2} \quad (11)$$

$$\Delta_i = \frac{U}{2} \langle c_i^\dagger \sigma_z c_i \rangle = (-1)^i \frac{UM}{2} = (-1)^i \Delta \quad (12)$$

where n is the average on-site electron density and cancels with the chemical potential μ in half-filled ($n = 1$) case. M is the staggered magnetization and $\Delta = \frac{UM}{2}$ is the energy band gap. The effective Lagrangian turns into

$$\mathcal{L}_{\text{eff}} = \sum_{i,\sigma} c_{i,\sigma}^* \partial_\tau c_{i,\sigma} - \sum_{\langle ij \rangle} (t_{i,j} c_i^* c_j + h.c.) - \sum_i (-1)^i \Delta c_i^* \sigma_z c_i. \quad (13)$$

One may obtain the spectrum of the electrons as

$$E_{\mathbf{k}} = \pm \sqrt{|\xi_{\mathbf{k}}|^2 + \Delta^2} \quad (14)$$

Finally we derive the self-consistency equation for M by minimizing the free energy at temperature T in the Brillouin zone as

$$1 = \frac{1}{N} \sum_{\mathbf{k}} \frac{U}{2E_{\mathbf{k}}} \tanh(\beta E_{\mathbf{k}}/2) \quad (15)$$

Here $\beta = \frac{1}{k_B T}$. N is the number of the sites.

From Fig.2, one could find that MI transition occurs at a critical value about $U/t \simeq 2.23$ at zero temperature^{14,15,16,23}. In the weakly coupling limit ($U/t < 2.23$), the ground state is a semi-metal (SM) with *nodal* fermi-points. In the strong coupling region ($U/t > 2.23$), due to $M \neq 0$, the ground state becomes




FIG. 2: Staggered magnetization M of half-filled honeycomb lattice Hubbard model at $T = 0$.

an insulator with massive fermionic excitations. However, the non-zero value of M only means the existence of effective spin moments rather than a long range AF order since this result is obtained in a mean field level by fixing the spins along $\hat{z}$ -axis. Thus one needs to examine stability of magnetic order against quantum fluctuations of effective spin moments by keeping spin rotation symmetry.

III. EFFECTIVE NONLINEAR σ MODEL IN THE INSULATOR STATE

In this section, we will derive an effective NL σ M of spin fluctuations with spin rotation symmetry in the honeycomb Hubbard model beyond above mean field theory.

To describe the spin fluctuations, we use the Haldane's mapping:

$$\mathbf{\Omega}_i = (-1)^i \mathbf{n}_i \sqrt{1 - \mathbf{L}_i^2} + \mathbf{L}_i \quad (16)$$

where $\mathbf{n}_i$ is the Neel vector that corresponds to the long-wavelength part of $\mathbf{\Omega}_i$ with a restriction $\mathbf{n}_i^2 = 1$. $\mathbf{L}_i$ is the transverse canting field that corresponds to the short-wavelength parts of $\mathbf{\Omega}_i$ with a restriction $\mathbf{L}_i \cdot \mathbf{n}_i = 0$ ^{18,24,25}. We then rotate $\mathbf{\Omega}_i$ to $\hat{z}$ -axis for the spin indexes of the electrons at i -site:^{17,18,19,20,21,22}

$$\begin{aligned} \psi_i &= U_i^\dagger c_i \\ U_i^\dagger \mathbf{n}_i \cdot \sigma U_i &= \sigma_z \\ U_i^\dagger \mathbf{L}_i \cdot \sigma U_i &= \mathbf{l}_i \cdot \sigma \end{aligned} \quad (17)$$

where $U_i \in \text{SU}(2)/\text{U}(1)$. One then can derive the follow-

ing effective Lagrangian after such spin transformation:

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= \sum_i \psi_i^* \partial_\tau \psi_i + \sum_i \psi_i^* a_0(i) \psi_i \\ &\quad - \sum_{\langle ij \rangle} (t_{i,j} \psi_i^* e^{ia_{ij}} \psi_j + h.c.) \\ &\quad - \Delta \sum_i \psi_i^* \left[(-1)^i \sigma_z \sqrt{1 - \mathbf{l}_i^2} + \mathbf{l}_i \cdot \sigma \right] \psi_i \end{aligned} \quad (18)$$

where the auxiliary gauge fields $a_{ij} = a_{ij,1} \sigma_x + a_{ij,2} \sigma_y$ and $a_0(i) = a_{0,1}(i) \sigma_x + a_{0,2}(i) \sigma_y$ are defined:

$$e^{ia_{ij}} = U_i^\dagger U_j, \quad a_0(i) = U_i^\dagger \partial_\tau U_i. \quad (19)$$

In terms of the mean field result $M = (-1)^i \langle \psi_i^* \sigma_z \psi_i \rangle$ as well as the approximations,

$$\sqrt{1 - \mathbf{l}_i^2} \simeq 1 - \frac{\mathbf{l}_i^2}{2}, \quad e^{ia_{ij}} \simeq 1 + ia_{ij},$$

we obtain the effective Hamiltonian as:

$$\begin{aligned} \mathcal{L}_{\text{eff}} &\simeq \sum_i \psi_i^* \partial_\tau \psi_i + \sum_i \psi_i^* [a_0(i) - \Delta \sigma \cdot \mathbf{l}_i] \psi_i \\ &\quad - \sum_{\langle ij \rangle} [t_{i,j} \psi_i^* (1 + ia_{ij}) \psi_j + h.c.] \\ &\quad - \Delta \sum_i (-1)^i \psi_i^* \sigma_z \psi_i + \Delta M \sum_i \frac{\mathbf{l}_i^2}{2}. \end{aligned} \quad (20)$$

By integrating out the fermion fields ψ_i^* and ψ_i , the effective action with the quadric terms of $[a_0(i) - \Delta \sigma \cdot \mathbf{l}_i]$ and a_{ij} becomes

$$\mathcal{S}_{\text{eff}} = \frac{1}{2} \int_0^\beta d\tau \sum_i [-4\varsigma (a_0(i) - \Delta \sigma \cdot \mathbf{l}_i)^2 + 4\rho_s a_{ij}^2 + \frac{2\Delta^2}{U} \mathbf{l}_i^2] \quad (21)$$

where ρ_s and ς are two parameters.

To learn the properties of the low energy physics, we study the continuum theory of the effective action in Eq.(21). In the continuum limit, we denote $\mathbf{n}_i, \mathbf{l}_i, ia_{ij} \simeq U_i^\dagger U_j - 1$ and $a_0(i) = U_i^\dagger \partial_\tau U_i$ by $\mathbf{n}(x, y), \mathbf{l}(x, y), U^\dagger \partial_x U$ (or $U^\dagger \partial_y U$) and $U^\dagger \partial_\tau U$, respectively. From the relations between $U^\dagger \partial_\mu U$ and $\partial_\mu \mathbf{n}$,

$$a_\tau^2 = a_{\tau,1}^2 + a_{\tau,2}^2 = -\frac{1}{4} (\partial_\tau \mathbf{n})^2, \quad \tau = 0, \quad (22)$$

$$a_\mu^2 = a_{\mu,1}^2 + a_{\mu,2}^2 = \frac{1}{4} (\partial_\mu \mathbf{n})^2, \quad \mu = x, y, \quad (23)$$

$$\mathbf{a}_0 \cdot \mathbf{l} = -\frac{i}{2} (\mathbf{n} \times \partial_\tau \mathbf{n}) \cdot \mathbf{l}, \quad (24)$$

the continuum formulation of the action in Eq.(21) turns into

$$\begin{aligned} \mathcal{S}_{\text{eff}} &= \frac{1}{2} \int_0^\beta d\tau \int d^2 \mathbf{r} [\varsigma (\partial_\tau \mathbf{n})^2 + \rho_s (\nabla \mathbf{n})^2 \\ &\quad - 4i\Delta \varsigma (\mathbf{n} \times \partial_\tau \mathbf{n}) \cdot \mathbf{l} + (\frac{2\Delta^2}{U} - 4\Delta^2 \varsigma) \mathbf{l}^2] \end{aligned} \quad (25)$$

where the vector $\mathbf{a}_0$ is defined as $\mathbf{a}_0 = (a_{0,1}, a_{0,2}, 0)$.

Finally we integrate the transverse canting field $\mathbf{l}$ and obtain the effective NLSM as

$$\mathcal{S}_{\text{eff}} = \frac{1}{2g} \int_0^\beta d\tau \int d^2r \left[\frac{1}{c} (\partial_\tau \mathbf{n})^2 + c (\nabla \mathbf{n})^2 \right] \quad (26)$$

with a constraint $\mathbf{n}^2 = 1$. The coupling constant g and spin wave velocity c are defined as:

$$g = \frac{c}{\rho_s}, \quad c^2 = \frac{\rho_s}{\chi^\perp}, \quad \chi^\perp = \left(\frac{1}{\zeta} - 2U \right)^{-1}. \quad (27)$$

Here ρ_s is the spin stiffness,

$$\rho_s = \frac{1}{N} \sum_{\mathbf{k}} \frac{\epsilon^2}{4(|\xi_{\mathbf{k}}|^2 + \Delta^2)^{\frac{3}{2}}}, \quad (28)$$

where the corresponding coefficient ϵ^2 is

$$\begin{aligned} \epsilon^2 = & \frac{1}{4} t^2 [6\Delta^2 + 27t^2 + (2\Delta^2 + 27t^2) \cos(\sqrt{3}k_y) \\ & + 36t^2 \cos(3k_x/2) \cos(\sqrt{3}k_y/2) \cos(\sqrt{3}k_y) \\ & + 2(5\Delta^2 + 27t^2) \cos(3k_x/2) \cos(\sqrt{3}k_y/2) \\ & + 9t^2 \cos(3k_x) (1 + \cos(\sqrt{3}k_y))]. \end{aligned} \quad (29)$$

$\chi^\perp$ is the transverse spin susceptibility, of which ζ is

$$\zeta = \frac{1}{N} \sum_{\mathbf{k}} \frac{\Delta^2}{4(|\xi_{\mathbf{k}}|^2 + \Delta^2)^{\frac{3}{2}}}. \quad (30)$$

One can see detailed calculations of ρ_s and ζ in the appendix.

The numerical results of ρ_s and c are illustrated in Fig.3, where one can find that $c = 0.168901985t = 1.055637J$ in the strongly coupling limit ($U = 25t$) match the earlier results, $c = 1.06066J$ ($J = \frac{4t^2}{U}$), obtained from Heisenberg model²⁶.

In addition, we need to determine another important parameter - the cutoff Λ . On the one hand, the effective NLSM is valid within the energy scale of Mott gap, $2\Delta = UM$. On the other hand, the lattice constant is a natural cutoff. Thus the cutoff is defined as the following equation¹⁸

$$\Lambda = \min(1, \frac{2\Delta}{c}) \quad (31)$$

IV. MAGNETIC PROPERTIES OF THE INSULATOR STATE

In this section we will use the effective NLSM to study the magnetic properties of the insulator state²⁷. The




FIG. 3: The spin stiffness ρ_s and the spin wave velocity c of the half-filled honeycomb Hubbard model at $T = 0$.




FIG. 4: The dimensionless coupling constant α of Hubbard model on honeycomb lattice of half filling at $T = 0$. Insert (b) shows non-magnetic region.

Lagrangian of NLSM with a constraint ($\mathbf{n}^2 = 1$) by a Lagrange multiplier λ becomes

$$\mathcal{L}_{\text{eff}} = \frac{1}{2cg} \left[(\partial_\tau \mathbf{n})^2 + c^2 (\nabla \mathbf{n})^2 + i\lambda(1 - \mathbf{n}^2) \right] \quad (32)$$

where $i\lambda = m^2$ and m is the mass gap of the spin fluctuations.

Using the large- N approximation we rescale the field $\mathbf{n} \rightarrow \sqrt{N}\mathbf{n}$ and obtain the saddle-point equation of motion as

$$(n_0)^2 - k_B T \sum_{\omega_n, \mathbf{q} \neq 0} \Pi(\mathbf{q}, i\omega_n) = 1. \quad (33)$$

In Eq.(33), n_0 is the mean field value of $\mathbf{n}$ and $\Pi(\mathbf{q}, i\omega_n) = -\frac{gc}{\omega_n^2 + c^2 \mathbf{q}^2 + m^2}$ is the propagator of the spin fluctuations $\delta \mathbf{n} = \mathbf{n} - n_0$. Here $\omega_n = 2\pi n k_B T$, $n = \text{integers}$.

At finite temperature, the solution of n_0 is always zero that is consistent to the Mermin-Wigner theorem. From




FIG. 5: The mass gap m of the spin fluctuations of Hubbard model on honeycomb lattice of half filling at $T = 0$.

Eq.(33), we may get the solution of m as

$$m = 2k_B T \sinh^{-1} \left[e^{-\frac{2\pi c}{g k_B T}} \sinh \left(\frac{c\Lambda}{2k_B T} \right) \right]. \quad (34)$$

In the limit $T \ll \Lambda$, Eq.(34) can be rewrite as:

$$m = 2k_B T \sinh^{-1} \left\{ \frac{1}{2} \exp \left[-\frac{2\pi c}{k_B T} \left(\frac{1}{g} - \frac{1}{g_c} \right) \right] \right\} \quad (35)$$

where

$$g_c = \frac{4\pi}{\Lambda}. \quad (36)$$

Therefore, at zero temperature the solutions of n_0 and m of Eq.(33) are determined by the dimensionless coupling constant $\alpha = g\Lambda$. In particular, there exists a critical point $\alpha_c = 4\pi$ (or $g_c = \frac{4\pi}{\Lambda}$): For the case of $\alpha < 4\pi$, we get solutions of n_0 and m :

$$n_0 = (1 - \frac{g}{g_c})^{1/2}, m = 0 \quad (37)$$

For the case of $\alpha > 4\pi$, we get solutions of n_0 and m :

$$n_0 = 0, m = 4\pi c \left(\frac{1}{g_c} - \frac{1}{g} \right) \quad (38)$$

So we calculate the dimensionless coupling constant $\alpha = g\Lambda$ and show results in Fig.4. The quantum critical points corresponding to $\alpha_c = 4\pi$ turn into $(U/t)_{c2} \simeq 2.88$ and $(U/t)_{c3} \simeq 2.93$ which divide the insulator state into three phases - a quantum disordered state (QD) in the region of $2.88 < U/t < 2.93$ and two long range AF order in the regions of $2.23 < U/t < 2.88$ and $U/t > 2.93$.

In the regions of $2.23 < U/t < 2.88$ and $U/t > 2.93$ (where $\alpha < \alpha_c$), at low temperature the mass gap m of spin fluctuations is determined by:

$$m \simeq 2k_B T \exp \left[-\frac{2\pi c}{k_B T} \left(\frac{1}{g} - \frac{1}{g_c} \right) \right] \quad (39)$$




FIG. 6: Spin order parameter $\mathcal{M}_0$ of the Hubbard model on honeycomb lattice of half filling at $T = 0$. Insert (b) shows non-magnetic region.

Because the energy scale of the mass gap m is always much smaller than the temperature, *i.e.*, $m \ll k_B T$ (or ω_n), quantum fluctuations become negligible in a sufficiently long wavelength and low energy regime ($m < |c\mathbf{q}| < k_B T$). Thus in this region one may only consider the purely static (semiclassical) fluctuations.

At zero temperature, the mass gap m vanishes (See Fig.5), which means that long range AF order appears. To describe the AF order, we introduce a spin order parameter^{28,29,30}:

$$\mathcal{M}_0 = \frac{M}{2} n_0 = \frac{M}{2} (1 - \frac{g}{g_c})^{1/2} \quad (40)$$

As shown in Fig.6, the ground state of AF ordered phase has a finite spin order parameter.

In the region of $2.88 < U/t < 2.93$ (where $\alpha > \alpha_c$), there is a finite mass gap of spin fluctuations at zero temperature (See Fig.5):

$$m = 4\pi c \left(\frac{1}{g_c} - \frac{1}{g} \right) \quad (41)$$

Therefore, the ground state of the insulator in this region is not a long range AF order. Instead, it is a quantum disordered state (or non-magnetic insulator state) with zero spin order parameter $\mathcal{M}_0 = 0$ in a *narrow non-magnetic window* (See Fig.6).

Based on above results, we get the global phase diagram which is illustrated in Fig.7. One can see that at finite T , there are four crossover lines, T_{HF} , T_ρ , T_ν , that separate five regions.

The highest crossover line is T_{HF} that is obtained from Eq.(15) and denotes the establish of the effective spin-moments. Above T_{HF} , it is metal phase without energy gap $\Delta = 0$. The crossover line $T_\rho \sim \rho_s$ denotes the validity of the NL σ M, where ρ_s is the energy scale of spin stiffness. In the region $T_\rho < T < T_{HF}$, the free spin-moments are established (denoted by $M \neq 0$) that show a Curie-Weiss behavior. In this region one




FIG. 7: Phase diagram of the Hubbard model on honeycomb lattice of half filling at finite temperature. Insert (b) shows non-magnetic region.

cannot use the effective $NL\sigma M$. Below T_p , short range spin-correlation exists and the effective $NL\sigma M$ is valid. The region below T_p is dominated by the crossover lines $T_v \sim \rho_s |1 - g/g_c|$ together with two quantum critical points (QCP) at $(U/t)_{c2} = 2.88$ and $(U/t)_{c3} = 2.93$ which represent the QCP of $g = g_c$ in the $NL\sigma M$.^{28,30}

V. SPIN-SPIN CORRELATIONS

In order to make our theoretical predictions verifiable, we discuss the detection of the anomalous spin dynamics in ultracold atom experiments via spatial spin-spin correlations $\langle \mathbf{S}(\mathbf{r}, t) \cdot \mathbf{S}(0, 0) \rangle = \langle c^\dagger(\mathbf{r}, t) \sigma c(\mathbf{r}, t) \cdot c^\dagger(0, 0) \sigma c(0, 0) \rangle$ or dynamic spin susceptibility $\chi''(\mathbf{q}, \omega)$. Here $\chi''(\mathbf{q}, \omega)$ is defined as

$$\chi''(\mathbf{q}, \omega) = \frac{1}{2} (1 - e^{-\omega\beta}) \int dt d\mathbf{r} e^{i(\omega t - \mathbf{q} \cdot \mathbf{r})} \langle \mathbf{S}(\mathbf{r}, t) \cdot \mathbf{S}(0, 0) \rangle. \quad (42)$$

On the one hand, people may observe spatial spin-spin correlations $\langle \hat{S}_z(\mathbf{r}_1) \hat{S}_z(\mathbf{r}_2) \rangle$ by noise correlation spectroscopy^{31,32}. As demonstrated in Ref.³¹, using a probe laser beam which goes through the system, one can measure the phase shift or change of polarization of the outgoing beam $\langle \hat{X}_{\text{out}} \rangle \propto \langle \hat{M}_z \rangle$ to obtain the magnetization $\langle \hat{M}_z \rangle \propto \int d\mathbf{r} \phi(\mathbf{r}) \langle \hat{S}_z(\mathbf{r}) \rangle$, where $\phi(\mathbf{r})$ is the spatial intensity profile of the laser beam. The quantum noise $\langle \hat{X}_{\text{out}}^2 \rangle \propto \langle \hat{M}_z^2 \rangle \propto \int d\mathbf{r}_1 d\mathbf{r}_2 \phi(\mathbf{r}_1) \phi(\mathbf{r}_2) \langle \hat{S}_z(\mathbf{r}_1) \rangle \langle \hat{S}_z(\mathbf{r}_2) \rangle$ reveals the atomic correlations in the system. This method can directly examine the existence of 2D AF correlations.

On the other hand, people may also observe dynamic spin susceptibility $\chi''(\mathbf{q}, \omega)$ which links with spin-spin correlations by NMR measurement. Consequently, it determines the spin-spin correlation in the real space as $\cos(\mathbf{Q}_0 \cdot \mathbf{r}) e^{-|\mathbf{r}|/\xi}$, with the correlation length ξ and the AF wave-vector $\mathbf{Q}_0 = (\pm \frac{\pi}{a}, \pm \frac{\pi}{a})$. To show clearly the




FIG. 8: Correlation length ξ at $T = 0.02, 0.1, 0.2$. The unit of temperature is t .

anomalous spin dynamics, we calculate the spin-spin correlation $\langle \mathbf{S}(\mathbf{r}, t) \cdot \mathbf{S}(0, 0) \rangle$. The dynamic spin susceptibility at $\mathbf{Q}_0$ is determined by n fields

$$\chi(\mathbf{q}, \omega) = \langle n_a(\mathbf{q}, \omega) n_a(-\mathbf{q}, -\omega) \rangle = \frac{gc}{\mathbf{q}^2 + \omega^2 + m^2} \quad (43)$$

where we have introduced the momentum $\mathbf{q}$. The equal-time spin-spin correlation function is proportion to $e^{-|\mathbf{r}|/\xi}$ where the spin-correlation length ξ is therefore given by $\frac{1}{m}$. As Eq.(42) indicates, the spin dynamic structure factor as well as the dynamic spin susceptibility function may reflect the effective short-range magnetic correlation length ξ .

In Fig.8, we give the spin-correlation length defined as $\xi = \frac{1}{m}$ at $T = 0.02t, 0.1t, 0.2t$, respectively. From the Fig.8, taking $T = 0.02t$ as an example, one can see that the spin-correlation length increases quickly with increasing interaction U/t . However, the spin-correlation length doesn't increase monotonously with U/t - it will decrease and reach a minimum value near the MI transition $U/t \sim (U/t)_{c2} \sim (U/t)_{c3}$. The dip of the spin-correlation length will indicate the existence of the non-magnetic state near MI transition. When one increases the interaction further, the spin-correlation increases again and finally decreases in the strongly interacting limit due to $J \rightarrow 0$. For other cases with higher temperature, $T = 0.1t, 0.2t$, there exist similar dip structure of the spin-correlation length via U/t which means that people may observe the anomalous spin dynamics more easily in experiments.

VI. DOPING EFFECT

In this section we will leave from half-filling and study the hole doping cases. In case of hole-concentration $d = 1 - n$, the chemical potential μ is not $\frac{U}{2}$ any more. The




FIG. 9: Spin stiffness ρ_s of the Hubbard model on honeycomb lattice with hole-concentration $d = 0.0, 0.1, 0.2, 0.3, 0.4, 0.5$ at $T = 0$.

Hamiltonian becomes

$$\mathcal{H} = - \sum_{\langle ij \rangle} (t_{ij} c_i^* c_j + h.c.) - \sum_i \Delta_i c_i^* \mathbf{\Omega}_i \cdot \boldsymbol{\sigma} c_i + \sum_i \left(\frac{U n}{2} - \mu \right) c_i^* c_i. \quad (44)$$

At this case, we could obtain M , μ , ζ , ρ_s similarly with that of half-filling case in $T = 0$ as follows,

$$1 = \frac{1}{N} \sum_{E_{\mathbf{k}} < \mu} \frac{U}{2E_{\mathbf{k}}}, \quad 1 - d = \sum_{E_{\mathbf{k}} < \mu} 1, \quad (45)$$

$$\zeta = \frac{1}{N} \sum_{E_{\mathbf{k}} < \mu} \frac{\Delta^2}{4 \left(|\xi_{\mathbf{k}}|^2 + \Delta^2 \right)^{\frac{3}{2}}},$$

$$\rho_s = \frac{1}{N} \sum_{E_{\mathbf{k}} < \mu} \frac{\epsilon^2}{4 \left(|\xi_{\mathbf{k}}|^2 + \Delta^2 \right)^{\frac{3}{2}}}.$$

Given certain hole-concentration d , the ρ_s and c may be obtained. From comparison in Fig.9, one can see that spin stiffness ρ_s rises when the hole-concentration increases from $d = 0.1$ to $d = 0.5$, and it shows a different behavior in the large doping density from that of low doping density. From Fig.10, the spin wave velocity c goes up as well when we increase hole concentration and it becomes much larger at critical points. The dimensionless coupling constant α are calculated in Fig.11 in different hole-concentrations, from which one can see that all critical points go up as we increase doping concentration d . And more importantly, we find that the non-magnetic regions widen when the hole-concentration increases.

In order to show the doping effect clearly, we plot the phase diagram of the honeycomb lattice Hubbard model at $T = 0$ (See Fig.12). From the phase diagram, one can see that there are three critical lines (U_{c1} , U_{c2} , U_{c3}) separating four regimes - one SM regime, two AF regimes




FIG. 10: Spin wave velocity c of the honeycomb lattice Hubbard model with hole-concentration $d = 0.0, 0.1, 0.2, 0.3, 0.4, 0.5$ at $T = 0$.




FIG. 11: The dimensionless coupling constant α of the Hubbard model on honeycomb lattice with hole-concentration $d = 0.0, 0.1, 0.2, 0.3, 0.4, 0.5$ at $T = 0$.

and one non-magnetic regime. In Fig.12, U_{c1} that has been given in Ref.²³ is derived by Hartree-Fock approximation. U_{c2} and U_{c3} are determined by the critical point of the effective NL σ M. In particular, the region of quantum disordered state becomes much wider by increasing the hole-concentration.

VII. CONCLUSION

In this paper, we investigate the two dimensional honeycomb Hubbard model with an approach that keeps spin rotation symmetry. By it, we found anomalous spin dynamics not far from the critical point of MI transition : there may exist a narrow non-magnetic insulator. The narrow non-magnetic insulator will be in favor of hole-doping. Also we discuss how to observe the anomalous spin dynamics in cold atoms.

In the end we discuss the nature of the non-magnetic




FIG. 12: Phase diagram of the Hubbard model on honeycomb lattice at different hole-concentration at $T = 0$. The square dot line, circle dot line, triangle dot line represent U_{c1} , U_{c2} , U_{c3} , respectively.

insulator in the two dimensional honeycomb Hubbard model. In Ref.³³, it is pointed out that by considering the fermionic nature of vortices (half-skyrmions), nodal spin liquid becomes the ground state of the non-magnetic insulator state in the π -flux Hubbard model on square lattice. There exist three types of quasi-particles in nodal spin liquids: nodal fermionic spinons, gapped bosonic spinons and roton-like $U(1)$ gauge field. Following the similar approach, one may draw the same conclusion that the non-magnetic insulator in the two dimensional honeycomb Hubbard model is another example nodal spin liquid with similar quasi-particles - nodal fermionic spinons, gapped bosonic spinons and roton-like $U(1)$ gauge field.

Acknowledgments

The authors acknowledge stimulating discussions with Z. Y. Weng, T. Li, N. H. Tong, F. Yang, P. Ye. This research is supported by NCET, NFSC Grant no. 10574014, 10874017.

APPENDIX A: THE DETAILED CALCULATIONS OF ρ_s AND ς

To give ρ_s and ς for calculation, we choose U_i to be

$$U_i = \begin{pmatrix} z_{i\uparrow}^* & z_{i\downarrow}^* \\ -z_{i\downarrow} & z_{i\uparrow} \end{pmatrix}, \quad (\text{A1})$$

where $\mathbf{n}_i = \bar{\mathbf{z}}_i \sigma \mathbf{z}_i$, $\mathbf{z}_i = (z_{i\uparrow}, z_{i\downarrow})^T$, $\bar{\mathbf{z}}_i \mathbf{z}_i = \mathbf{1}$. And the spin fluctuations around $\mathbf{n}_i = \hat{\mathbf{z}}_i$ is

$$\mathbf{n}_i = \hat{\mathbf{z}}_i + \text{Re}(\phi_i) \hat{\mathbf{x}} + \text{Im}(\phi_i) \hat{\mathbf{y}} \quad (\text{A2})$$

$$\mathbf{z}_i = \begin{pmatrix} 1 - |\phi_i|^2/8 \\ \phi_i/2 \end{pmatrix} + O(\phi_i^3). \quad (\text{A3})$$

Then the quantities $U_i^\dagger U_j$ and $U_i^\dagger \partial_\tau U_i$ can be expanded in the power of $\phi_i - \phi_j$ and $\partial_\tau \phi_i$,

$$U_i^\dagger U_j = e^{i \frac{\phi_i - \phi_j}{2} \sigma_y} \quad (\text{A4})$$

$$U_i^\dagger \partial_\tau U_i = \begin{pmatrix} 0 & -\frac{1}{2} \partial_\tau \phi_i \\ \frac{1}{2} \partial_\tau \phi_i & 0 \end{pmatrix}. \quad (\text{A5})$$

According to Eq.(19), the gauge field a_{ij} and $a_0(i)$ are given as

$$a_{ij} = \frac{1}{2} (\phi_i - \phi_j) \sigma_y \quad (\text{A6})$$

$$a_0(i) = \frac{i}{2} \partial_\tau \phi_i \sigma_y. \quad (\text{A7})$$

Supposing a_{ij} and $a_0(i)$ to be a constant in space and denoting $\partial_i \phi_i = \mathbf{a}$ and $\partial_\tau \phi_i = i B_y$, we have

$$a_{ij} = -\frac{1}{2} \mathbf{a} \cdot (\mathbf{i} - \mathbf{j}) \sigma_y \quad (\text{A8})$$

$$a_0(i) = -\frac{1}{2} B_y \sigma_y. \quad (\text{A9})$$

The energy of Hamiltonian of Eq.(21) becomes

$$E(B_y, \mathbf{a}) = \frac{1}{2} \zeta B_y^2 + \frac{1}{2} \rho_s \mathbf{a}^2. \quad (\text{A10})$$

Then one could get ζ and ρ_s from the following equations by calculating the partial derivative of the energy

$$\zeta = \frac{1}{N} \frac{\partial^2 E_0(B_y)}{\partial B_y^2} \Big|_{B_y=0} \quad (\text{A11})$$

$$\rho_s = \frac{1}{N} \frac{\partial^2 E_0(\mathbf{a})}{\partial \mathbf{a}^2} \Big|_{\mathbf{a}=0}. \quad (\text{A12})$$

Here $E_0(B_y)$ and $E_0(\mathbf{a})$ are the energy of the lower Hubbard band

$$E_0(B_y) = \sum_{\mathbf{k}} (E_{+, \mathbf{k}}^\zeta + E_{-, \mathbf{k}}^\zeta) \quad (\text{A13})$$

$$E_0(\mathbf{a}) = \sum_{\mathbf{k}} (E_{+, \mathbf{k}}^\rho + E_{-, \mathbf{k}}^\rho) \quad (\text{A14})$$

where $E_{+, \mathbf{k}}^\zeta$, $E_{-, \mathbf{k}}^\zeta$ and $E_{+, \mathbf{k}}^\rho$, $E_{-, \mathbf{k}}^\rho$ are the energies of the following Hamiltonian $\mathcal{H}^\zeta$ and $\mathcal{H}^\rho$

$$\begin{aligned} \mathcal{H}^\zeta = & - \sum_{\langle ij \rangle} (t_{i,j} \psi_i^* \psi_j + h.c.) - \Delta \sum_i (-1)^i \psi_i^* \sigma_z \psi_i \\ & + \sum_i \psi_i^* a_0(i) \psi_i \end{aligned} \quad (\text{A15})$$

$$\mathcal{H}^\rho = - \sum_{\langle ij \rangle} (t_{i,j} \psi_i^* e^{a_{ij}} \psi_j + h.c.) - \Delta \sum_i (-1)^i \psi_i^* \sigma_z \psi_i. \quad (\text{A16})$$

Using the Fourier transformations for $\mathcal{H}^\zeta$, we have the spectrum of the lower band of $\mathcal{H}^\zeta$

$$E_{\pm, \mathbf{k}}^\zeta = -\sqrt{\left(|\xi_{\mathbf{k}}| \pm \frac{B_y}{2} \right)^2 + \Delta^2} \quad (\text{A17})$$

where $\xi_{\mathbf{k}}$ has been obtained in Eq.(6). Using Eq.(A11), ζ is obtained as

$$\zeta = \frac{1}{N} \sum_{\mathbf{k}} \frac{\Delta^2}{4(|\xi_{\mathbf{k}}|^2 + \Delta^2)^{\frac{3}{2}}}. \quad (\text{A18})$$

Similarly, using the Fourier transformations for $\mathcal{H}^\rho$, we obtain the spectrum of the lower band of $\mathcal{H}^\rho$

$$E_{\pm, \mathbf{k}}^\rho = -\sqrt{\Delta^2 + |\psi|^2 + |\varphi|^2 \pm [4\Delta^2 |\psi|^2 - (\varphi\psi^* - \psi\varphi^*)^2]^{\frac{1}{2}}} \quad (\text{A19})$$

where φ and ψ are defined as

$$\varphi = -t \sum_{\delta} e^{i\mathbf{k} \cdot \delta} \cos\left(\frac{1}{2}\mathbf{a} \cdot \delta\right) \quad (\text{A20})$$

$$\psi = -t \sum_{\delta} e^{i\mathbf{k} \cdot \delta} \sin\left(\frac{1}{2}\mathbf{a} \cdot \delta\right). \quad (\text{A21})$$

Using Eq.(A12), ρ_s is given as

$$\rho_s = \frac{1}{N} \sum_{\mathbf{k}} \frac{\epsilon^2}{4(|\xi_{\mathbf{k}}|^2 + \Delta^2)^{\frac{3}{2}}}. \quad (\text{A22})$$

The corresponding coefficient ϵ^2 is

$$\begin{aligned} \epsilon^2 = & \frac{1}{4} t^2 [6\Delta^2 + 27t^2 + (2\Delta^2 + 27t^2) \cos(\sqrt{3}k_y) \\ & + 36t^2 \cos(3k_x/2) \cos(\sqrt{3}k_y/2) \cos(\sqrt{3}k_y) \\ & + 2(5\Delta^2 + 27t^2) \cos(3k_x/2) \cos(\sqrt{3}k_y/2) \\ & + 9t^2 \cos(3k_x) (1 + \cos(\sqrt{3}k_y))]. \end{aligned} \quad (\text{A23})$$

-
- * Corresponding author; Electronic address: spkou@bnu.edu.cn
- ¹ I. Bloch, J. Dalibard and W. Zwerger *Rev. Mod. Phys.* **80** 885 (2008).
 - ² S. Giorgini, L. P. Pitaevskii, & S. Stringari, *Rev. Mod. Phys.* **80**, 1215 (2008).
 - ³ M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, & I. Bloch, *Nature* **415**, 39-44 (2002).
 - ⁴ R. Jödens, N. Strohmaier, K. Günter, H. Moritz, & T. A. Esslinger, *Nature* **455**, 204-207 (2008).
 - ⁵ U. Schneider, *Science* **322**, 1520-1525 (2008).
 - ⁶ M. Iskin and C. J. Williams, *Phys. Rev. A* **78**, 011603(R) (2008).
 - ⁷ B. M. Andersen and G. M. Bruun, *Phys. Rev. A* **76**, 041602(R) (2007).
 - ⁸ M. Snoek, I. Titvinidze, C. Toke, K. Byczuk, and W. Hofstetter, *New J. Phys.* **10**, 093008 (2008).
 - ⁹ T. Gottwald and P. G. J. van Dongen, *Phys. Rev. A* **80**, 033603 (2009).
 - ¹⁰ L.-M. Duan, E. Demler, and M. D. Lukin, *Phys. Rev. Lett.* **91**, 090402 (2003).
 - ¹¹ S. L. Zhu, B. G. Wang, and L.-M. Duan, *Phys. Rev. Lett.* **98**, 260402 (2007).
 - ¹² Kean Loon Lee *et al.*, *A* **80**, 043411 (2009).
 - ¹³ M. Hermele, *Phys. Rev. B*, **76** 035125 (2007).
 - ¹⁴ S. Sorella and E. Tosatti, *Europhys. Lett.* **19**, 699 (1992).
 - ¹⁵ L. M. Martelo, *et al.*, *Z. Phys. B: Condens. Matter* **103**, 335 (1997).
 - ¹⁶ T. Paiva, *et al.*, *Phys. Rev. B* **72**, 085123 (2005).
 - ¹⁷ N. Dupuis, *Phys. Rev. B* **65**, 245118 (2002).
 - ¹⁸ K. Borejsza, N. Dupuis, *Euro Phys. Lett.* **63**, 722 (2003).
K. Borejsza and N. Dupuis *Phys. Rev. B* **69**, 085119 (2004).
 - ¹⁹ G. Y. Sun and S. P. Kou, *Europhys. Lett.* **87** 67002 (2009).
 - ²⁰ X. G. Wen, *Quantum Field Theory of Many-Body Systems*, (Oxford Univ. Press, Oxford, 2004).
 - ²¹ H. J. Schulz, in *The Hubbard Model*, edited by D. Baeriswyl (Plenum, New York, 1995).
 - ²² Z. Y. Weng, C. S. Ting, and T. K. Lee, *Phys. Rev. B* **43**, 3790 (1991).
 - ²³ N. M. R. Peres, *et al.*, *Phys. Rev. B* **70**, 195122 (2004).
 - ²⁴ F. D. M. Haldane, *Phys. Lett.* **93A**, 464 (1983).
 - ²⁵ A. Auerbach, *Interacting Electrons and Quantum Magnetism* (Springer-Verlag, New York, 1994).
 - ²⁶ Ann Mattsson and Per Fröjdh, *Phys. Rev. B* **49**, 3997 (1994).
 - ²⁷ Near the critical point that the mean field value of M vanishes, due to the strong charge fluctuations and the strong amplitude fluctuations of M , NL σ M cannot be used and our results cannot be reliable.
 - ²⁸ S. Chakravarty, *et al.*, *Phys. Rev. B* **39**, 2344 (1989).
 - ²⁹ A.V. Chubukov and D.M. Frenkel, *Phys. Rev. B* **46**, 11884 (1992).
 - ³⁰ S. Sachdev, *Quantum Phase Transitions*, (Cambridge University Press, 1999).
 - ³¹ G. M. Bruun, B. M. Andersen, E. Demler, A. S. Sørensen, *Phys. Rev. Lett.* **102**, 030401 (2009).
 - ³² V. Guarrera, *et al.*, *Phys. Rev. Lett.* **100**, 250403 (2008).
 - ³³ S. P. Kou, L. F. Liu, arXiv: 0910.2070.